
Yakima County Population =217,700 (31% under age 18)

Adult Tobacco Use

Current Cigarette Smoking (1987-1998)




Current Smokeless Tobacco Use (1993-1997)

Yakima County Washington State Entire adult population 3.5% (95%CI:2.0,6.2) 2.9% (95%CI:2.5,3.8)

Males only 6.7% (95%CI:3.7,12.0) 5.7% (95%CI:5.0,6.5)

Youth Tobacco Use (1998)



Summary of the problem

- ☐ Yakima County and state adult tobacco use rates are not significantly different.
- Youth tobacco use for the Eastern Washington region is elevated above the state rate for both cigarettes and smokeless tobacco, among all age groups. More data collection is needed to describe youth tobacco use in Yakima County.
- ☐ Maternal smoking during pregnancy has been consistently and significantly lower in Yakima than the state since 1990.
- Retailer compliance seems to be worse in Yakima County than in the state (retailers more likely to sell tobacco to minors).




Smoking During Pregnancy (1987-1998)

30% Yakima County Washington % smoking during pregnancy 25% 20% 15% 10% 5% 0% 1885 1003

Minors Access to Tobacco (1998-1999)

Percent Sales to Minors



Infants born annually to mothers who smoke during pregnancy: 456

Number of tobacco retailers in county: 322

Health at Risk (Estimated numbers for county)

Adults who currently smoke	38,100	
Adults who currently use smokeless tobacco	4,400	
Youth who currently smoke	4,700	
Youth who currently use smokeless tobacco	2,000	
ESTIMATED ANNUAL DEATHS DUE TO TOBACCO USE	340	

Current Tobacco Prevention and Control Program Activities Highlights during 1999

- Trained 77 Teens As Teachers peer educators from 12 county high schools and one middle school. Provided tobacco presentations to over 500 younger students.
- November 18, 1999: Approximately 600 students attended the multi-agency sponsored Burial of the Marl Burro Man public awareness event, held in conjunction with the Great American Smokeout and as the kick-off event for statewide youth coalition.
- Completed 38 tobacco retailer compliance checks.

Annual Funding for Program (12-month funding amounts, FY 1999)

Source	Amount	Agency receiving funding
Youth Access (Retailer License Fees)	\$ 24,217	American Lung Association
CDC Grant	\$ 0	•
Total Per Capita Funding	\$.11	CDC Recommended Community Per Capita Funding: \$.70-2.00

Local Contacts

Hailey Croci/Leslie Benoit American Lung Association 110 South 9th Avenue Yakima, WA 98902 (509) 248-4384 fax: (509) 248-4943

email: alawyakima1@nwinfo.net

Website: www.alaw.org



TECHNICAL NOTES: (ADDITIONAL DOCUMENTATION IN APPENDIX B)

TOTAL DEATHS DUE TO TOBACCO USE ARE ESTIMATED AS 19% OF ALL DEATHS (MCGINNIS AND FOEGE 1993); "DUMBBELL" BARS AROUND DATA POINTS FOR SURVEY DATA REPRESENT 95% CONFIDENCE INTERVALS - THE TRUE VALUE MAY BE ANYWHERE WITHIN THE BARS, BUT MAY NOT BE EXACTLY AT THE DATA POINT; NUMBER OF ADULTS/YOUTH WHO WILL DIE PREMATURELY FROM TOBACCO USE IS ESTIMATED AS 33% OF CURRENT ADULT SMOKERS (MMWR 1996), AND 21% OF CURRENT YOUTH SMOKERS -WHO MAY 'GROW OUT OF' SMOKING BEHAVIOR (USDHHS 1994): "CURRENT ADULT/YOUTH TOBACCO USERS" ARE SYNTHETIC ESTIMATES BASED ON STATEWIDE AGE-SPECIFIC PREVALENCE: FUNDING LEVELS FOR COUNTIES WERE CALCULATED BASED. ON DEDICATED YOUTH TOBACCO PREVENTION FUNDS AND CDC FUNDS MANAGED BY THE DEPARTMENT OF HEALTH - COUNTIES MAY SUCCESSFULLY SEEK ADDITIONAL FUNDING FOR TOBACCO CONTROL FROM OTHER SOURCES; "HIGHLIGHTS DURING 1999" WERE IDENTIFIED BY LOCAL PROGRAM STAFF; "CDC RECOMMENDED FUNDING" IS BASED ON BEST PRACTICES FOR TOBACCO CONTROL PROGRAMS (CDC 1999).